

A GENERALIZATION OF LAYMAN–LOTOCKII EXPONENTIAL INTERPOLATION

HERBERT E. SALZER

941 Washington Avenue Brooklyn NY 11225 U.S.A.

(Received June 1982; in revised form May 1983)

Communicated by Ervin Y. Rodin

Abstract—For equally spaced data points $f(i)$, $i = 0(1)n$, an exponential interpolation polynomial for $f(z)$, namely $\sum_{i=0}^n a_i r_i^z$, where r_i are arbitrary quantities, is obtainable formally from the Newton divided difference series for the displacement operator $E^z = (1 + \Delta)^z$, regarding Δ as the variable, for $\Delta = s_i = r_i - 1$. This generalization of Layman–Lotockiï exponential interpolation in which $r_i = i + 1$, employs “general factorial differences” $\Pi_{j=0}^i (\Delta - s_j) f(0)$, $i = 0(1) n - 1$. For the confluent case, where $f^{(i)}(0)$ is given instead of $f(i)$, $i = 0(1)n$, the results are similar, the derivative operator D replacing the Δ .

INTRODUCTION

Layman[1–4] and Lotockiï[5], working apparently independently around the same time, studied this interpolation series for $f(z)$ given at equally spaced arguments which are transformable into 0, 1, 2, . . . ,

$$f(z) \sim \sum_{i=0}^{\infty} \binom{\Delta}{i} f(0) e_i(z), \quad (1)$$

where $\binom{\Delta}{i} f(0)$ is proportional to the i th “factorial difference” $\Delta(\Delta - 1) \dots (\Delta - i + 1) f(0)$, and where

$$e_i(z) = \sum_{j=0}^i (-1)^j \binom{i}{j} (i + 1 - j)^z, \quad (2)$$

is the i th advancing difference at $\Delta = 0$ of the displacement operator $E^z = (1 + \Delta)^z$ in which Δ , instead of z , is regarded as the variable. From (1) and (2) we obtain for $f(z)$ an $(n + 1)$ -point exponential interpolation polynomial of the form $\sum_{i=0}^n a_i (i + 1)^z$.

We consider here the more general problem of finding the series corresponding to (1), (2), for equally spaced data points $f(i)$, $i = 0, 1, 2, \dots$, which yields an $(n + 1)$ -point exponential interpolation polynomial of the form $\sum_{i=0}^n a_i r_i^z$ in which the r_i 's are arbitrary quantities, not necessarily real, even though $r_i > 0$ is usually specified when $f(i)$ is real. Layman saw nothing better than Cramer's rule for finding $\sum_{i=0}^n a_i r_i^z$ ([1], p. 118). We show here, by the appropriate expansion of the operator $(1 + \Delta)^z$, how to produce the series which generalizes (1), (2), and from which we obtain $\sum_{i=0}^n a_i r_i^z$ directly and much more expediently than by Cramer's rule.

We also note briefly the confluent case, where we are given $f^{(i)}(0)$ instead of $f(i)$, $i = 0, 1, \dots$, from which we derive similar exponential series involving the derivative operator D instead of Δ , from e^{Dz} in place of $(1 + \Delta)^z$.

Convergence of the interpolation series (3), and (10) below, are still open problems, approaches to which might be suggested by results of Layman and Lotockiï concerning the convergence of $\sum_{i=0}^{\infty} a_i e_i(z)/i!$ for a given sequence of a_i 's and the representability of a given entire function $f(z)$ by the series in (1)[3–5]. For example, in connection with representability there is Layman's theorem on the convergence of (1) for $f(z)$ an entire function of exponential type, provided that a certain set $D(f)$ (see [3], p. 519 for a full description) lies within some suitably defined strip of width π ([3], p. 521).

INTERPOLATION SERIES

Regarding the operator $(1 + \Delta)^z$ as a function of Δ , with z as a parameter, expand it as a Newton divided difference series, for $\Delta = s_i = r_i - 1$, $i = 0, 1, 2, \dots$, and apply it to $f(z)$

at $z = 0$, obtaining this formal infinite series:

$$\begin{aligned} f(z) = E^z f(0) = (1 + \Delta)^z f(0) &= [s_0] f(0) + [s_0 s_1] (\Delta - s_0) f(0) + [s_0 s_1 s_2] (\Delta - s_0) (\Delta - s_1) f(0) \\ &+ \cdots + [s_0 s_1 \dots s_i] (\Delta - s_0) (\Delta - s_1) \dots (\Delta - s_{i-1}) f(0) \\ &+ \cdots, \end{aligned} \quad (3)$$

in which the divided differences $[s_0] = r_0^z$ and

$$[s_0 s_1 \dots s_i] = \sum_{j=0}^i r_j^z / \Pi_{k=0, k \neq j}^i (s_j - s_k). \quad (4)$$

The operator $\Pi_{k=0}^{i-1} (\Delta - s_k)$ may be defined as the “ i th general factorial difference operator”. Notice that

$$\left\{ \Pi_{k=0}^n (\Delta - s_k) \right\} \sum_{i=0}^n a_i r_i^z \equiv 0, \quad (5)$$

for all values of z , and Δ for a unit interval in z , i.e. whenever $f(z) \equiv \sum_{i=0}^n a_i r_i^z$, its $(n+1)$ th general factorial difference vanishes identically. The proof of (5) is immediate from the annihilation of $a_i r_i^z$ by the operational factor $\Delta - s_k$. Thus (5) is analogous to the vanishing of the $(n+1)$ th difference of any n th degree polynomial, and it may be employed in a similar checking capacity.

When $f(z) = \sum_{i=0}^n a_i r_i^z$ and the r_i 's are not known initially, they can be found from just $2n+2$ data points by the well-known Prony method.

When $z = n$ is within a region of convergence to $f(z)$ of the infinite series in (3), which breaks off after the $(n+1)$ th term when $z = n$ (since $[s_0 s_1 \dots s_i]$ in (4) is then the i th divided difference of the polynomial $(1 + \Delta)^n$, which vanishes for any choice of s_0, s_1, \dots, s_i , whenever $i > n$), the right member of (3) gives $f(n)$ for $z = n$. However, even when the infinite series in (3) may not converge to $f(z)$, it is still true that the first $n+1$ terms gives an exponential polynomial $\sum_{i=0}^n a_i r_i^z$ which equals $f(i)$ when $z = i$, $i = 0(1)n$ (fundamental interpolation property). This follows because, for $z = i$, (a) the $(i+2)$ th through the $(n+1)$ th terms of (3) vanish, being proportional to the $(i+j)$ th divided difference $[s_0 s_1 \dots s_{i+j}]$, $j = 1(1)n - i$, of $(1 + \Delta)^i$, and (b) the first $i+1$ terms are the result of an operator in Δ of the i th degree acting on $f(0)$, that operator (being any of the possible divided difference expansions in Δ -as-variable, of the i th degree polynomial $(1 + \Delta)^i$, which makes it equal to $(1 + \Delta)^i$ for all numerical values of Δ), having to be identical with $(1 + \Delta)^i$ as a polynomial function of Δ , from which we have $(1 + \Delta)^i f(0) = E^i f(0) = f(i)$. q.e.d. There is a slight subtlety in the preceding argument that goes from operator to polynomial and finally back to operator.

The uniqueness of the $(n+1)$ -point exponential interpolation polynomial $\sum_{i=0}^n a_i r_i^z$ follows from the non-vanishing $(n+1)$ th order determinant $\|r_i^j\|$, $i, j = 0(1)n$, which is a transposed Vandermondian. Thus the series (3), when truncated after the $(n+1)$ th term, reproduces any $f(z) \equiv \sum_{i=0}^n a_i r_i^z$ from its values at $z = 0(1)n$.

From (3) and (4) we obtain this explicit expression for the a_i in the interpolating $\sum_{i=0}^n a_i r_i^z$ (definitely preferable to Cramer's rule, when n is large):

$$a_i = \left\{ \sum_{j=i}^n [\Pi_{l=0}^{j-1} (\Delta - s_l) / \Pi_{l=0, l \neq i}^j (s_i - s_l)] \right\} f(0), \quad (6)$$

in which, for a_0 , for $j = 0$, $\Pi_{l=0}^{-1} (\Delta - s_l) / \Pi_{l=0, l \neq 0}^0 (s_0 - s_l) \equiv 1$.

To illustrate (6), we find the coefficients a_i , $i = 0(1)10$, in $f(z) = a_0 1.2^z + a_1 1.7^z + a_2 2.4^z + a_3 2.5^z + a_4 3^z + a_5 3.4^z + a_6 3.7^z + a_7 3.9^z + a_8 4.7^z + a_9 5.1^z + a_{10} 5.6^z$, having been given $f(z)$ to $10D$ for $z = 0(1)10$ (Table 1A, 1st column). The first ten general factorial differences, computed from the rounded $f(z)$'s (Table 1A, 2nd–11th columns) are all employed in obtaining the first row of Table 1A, the only one needed for (6). The size of $s_i = r_i - 1$, $i = 0(1)10$, increases the errors in successive columns, an *U.B.* for which was derived from the expansion of the operator $\Pi_{l=0}^{i-1} (\Delta - s_l)$. The denominators in (6), $\Pi_{l=0, l \neq i}^i (s_i - s_l)$ (Table 1B), magnify the errors when the r_i 's are close together. The a_i 's

found from (6) deviated from the true values of $a_i = (-1)^i + 3/(i+1)$ that were employed in the precomputation of $f(z)$, by somewhat more than 50% of the *U.B.* Considering that the system of 11 linear equations for the a_i 's is not too well-conditioned, the results, shown in Table 1C, might be regarded as satisfactory.

An equivalent alternative form for a_i , which is obtainable directly from the Lagrange interpolation polynomial form of the operator $(1 + \Delta)^z$, is not recommended because it requires $n+1$ sets of general factorial differences instead of just the single set in (6).

The interpolating $\sum_{i=0}^n a_i r_i^z$ is expressible also in this $(n+1)$ -point "Lagrangian exponential interpolation polynomial" form:

$$\sum_{i=0}^n a_i r_i^z = \sum_{i=0}^n L_i^{(n)}(z) f(i), \quad (7)$$

where, from (3) and (4), noting that $\Delta - s_k = E - r_k$ and that $f(i)$ first occurs in the $(i+1)$ th term of (3), we have

$$L_i^{(n)}(z) = \sum_{j=i}^n \sigma_{i,j} [s_0 \cdots s_j], \quad (8)$$

in which $\sigma_{i,j} = (-1)^{j-i} \times$ that elementary symmetric function of r_0, \dots, r_{j-1} which is of the $(j-i)$ th degree.

Table 1

1A. $f(z)$ and general factorial differences

z	$f(z)$	$(\Delta - s_0)f(z)$	$(\Delta - s_1)(\Delta - s_0)f(z)$
0	10.05963 20346	13.52559 52382	33.28235 71427
1	25.59715 36797	56.27586 90476	160.97162 14287
2	86.99245 34632	256.64059 88096	814.18676 92855
3	361.03154 29654	1250.47578 72618	4246.04555 33573
4	1683.71363 88203	6371.85439 17024	22609.40583 91213
5	8392.31075 82868	33441.55830 50154	1 22134.97448 78195
6	43512.33121 49596	1 78985.62360 63457	6 66508.71172 98481
7	2 31200.42106 42972	9 70784.27186 06359	36 64110.49860 64295
8	12 48224.77713 77925	53 14443.76076 95105	202 53280.98589 84190
9	68 12313.49333 48615	292 87835.37920 65868	
10	374 62611.57120 84206		
U.B. for error $\frac{1}{2} \times 10^{-10}$			
z	$(\Delta - s_2) \cdots (\Delta - s_0)f(z)$	$(\Delta - s_3) \cdots (\Delta - s_0)f(z)$	$(\Delta - s_4) \cdots (\Delta - s_0)f(z)$
0	81.09396 42862	225.11996 71411	547.00021 10073
1	427.85487 78566	1222.36011 24306	3021.82290 60917
2	2291.99730 70721	6688.90324 33835	16758.44946 61184
3	12418.89651 10638	36825.15919 62689	93228.29418 54536
4	67872.40047 39284	2 03703.77177 42603	5 19916.34273 43099
5	3 73384.77295 90813	11 31027.65805 70908	29 05108.83893 47305
6	20 64489.59045 47941	62 98191.81310 60029	
7	114 59415.78924 29882		
U.B. for error 10^{-9}			
z	$(\Delta - s_3) \cdots (\Delta - s_0)f(z)$	$(\Delta - s_6) \cdots (\Delta - s_0)f(z)$	$(\Delta - s_7) \cdots (\Delta - s_0)f(z)$
0	1162.02218 86669	2184.76948 73391	3737.23413 40241
1	6484.25158 54066	12257.83513 46466	21011.19127 62373
2	36249.56600 06510	68816.74830 13590	1 18129.42799 88362
3	2 02940.14250 37677	3 86514.74637 41363	
4	11 37393.27363 80768		
U.B. for error 6.22×10^{-8}			
z	$(\Delta - s_6) \cdots (\Delta - s_0)f(z)$	$(\Delta - s_9) \cdots (\Delta - s_0)f(z)$	
0	3446.19084 63240	1801.25568 42685	
1	19376.82900 05209		
U.B. for error 8.17×10^{-6}			
z	$(\Delta - s_9) \cdots (\Delta - s_0)f(z)$		
0			
1			

Table 1 (contd.)

1B. $\Pi'_{i=0, i \neq (s_i - s_j)}$		$j =$									
$i =$		1	2	3	4	5	6	7	8	9	10
0	-0.5	0.6	-0.78	1.404	-3.0888	7.722	-20.8494	72.9729	-284.59431	1252.214964	
1	0.5	-0.35	0.28	-0.364	0.6188	-1.2376	2.72272	-8.16816	27.771744	-108.3098016	
2		0.84	-0.084	0.0504	-0.0504	0.06552	-0.09828	0.226044	-0.6103188	1.95302016	
3			0.104	-0.052	0.0468	-0.05616	0.078624	-0.1729728	0.44972928	-1.394160768	
4				0.702	-0.2808	0.19656	-0.176904	0.3007368	-0.63154728	1.642022928	
5					1.3464	-0.40392	0.20196	-0.262548	0.4463316	-0.98192952	
6						1.638	-0.3276	0.3276	-0.45864	0.871416	
7							1.12266	-0.898128	1.0777536	-1.83218112	
8								93.93384	-37.573536	33.8161824	
9									223.315118208	-111.657559104	
10										1415.272345344	

1C. $a_i, i = 0(1)10$

i	0	1	2	3	4
a_i from (6)	3.99999 99341	0.50000 062	1.99997 24	-0.24996 04	1.60001 41
true value	4	0.5	2	-0.25	1.6
error	6.59×10^{-8}	6.2×10^{-7}	2.76×10^{-5}	3.96×10^{-5}	1.41×10^{-5}
<hr/>					
5	6	7	8	9	10
-0.49995 74	1.428526	-0.62497 947	1.33333 2376	-0.69999 729	1.27272 253
-0.5	1.428571...	-0.625	1.33333 3333...	0.7	1.27272 2727...
4.26×10^{-5}	4.5×10^{-5}	2.05×10^{-5}	10^{-6}	2.71×10^{-7}	2×10^{-8}

FACTORIAL DERIVATIVE SERIES

Confluent exponential interpolation series, similar to (3), (4) but employing derivatives D^j instead of differences, are derivable formally from the expansion of the operator $E^z = e^{Dz}$, regarding D as the variable. This applies when, in place of equally spaced data points, we have $D^j f(0) \equiv f^{(j)}(0)$, $j = 0, 1, 2, \dots$

The Gregory-Newton expansion of e^{Dz} , for $D = 0, h, 2h, \dots$ gives

$$f(z) = e^{Dz} f(0) = \left\{ 1 + \left(\frac{e^{hz} - 1}{h} \right) D + \left(\frac{e^{hz} - 1}{h} \right)^2 D(D-h)/2 + \dots + \left(\frac{e^{hz} - 1}{h} \right)^i D(D-h) \dots (D-(i-1)h)/i! + \dots \right\} f(0), \quad (9)$$

which converts a power series in z to one in $[(e^{hz} - 1)/h]$, with coefficients proportional to the "factorial derivatives" $D(D-h) \dots (D-(i-1)h)f(0)$. The series in (9) is the confluent form of the Layman-Lotockii series (1), (2).

The following more novel series, which is a generalization of (9), is obtained from the divided difference expansion of the operator e^{Dz} for $D = t_0, t_1, \dots$:

$$f(z) = e^{Dz} f(0) = \left\{ e^{t_0 z} + [(e^{t_1 z} - e^{t_0 z})/(t_1 - t_0)](D - t_0) + \dots + \left[\sum_{j=0}^i e^{t_j z} / \prod_{k=0, k \neq j}^i (t_j - t_k) \right] (D - t_0) \dots (D - t_{i-1}) + \dots \right\} f(0), \quad (10)$$

converting a power series into an exponential series, its coefficients being linear combinations of "general factorial derivatives" $(D - t_0) \dots (D - t_{i-1})f(0)$. For $t_j = \log r_j$, (10) is a series in r_j^z , of the same form as (3), (4), in which D and $\log r_j$ replace the Δ and $s_j = r_j - 1$.

That the first $n+1$ terms of (10) (this includes (9) as a special case), with their i th derivatives, at $z = 0$, is equal to $f^{(i)}(0)$, $i = 0(1)n$, follows from the quantities in brackets [...] becoming the successive divided differences of D^i considered as a quantity, for $D = t_j$, $j = 0(1)n$ (the $(i+1)$ th bracket, and those after, vanishing), from which the expression within the braces $\{\dots\}$, becoming the $(i+1)$ -point Newton divided difference identity for the polynomial D^i , must reproduce D^i considered as an operator. q.e.d.

The uniqueness of the $(n+1)$ -point completely osculatory exponential sum $\sum_{j=0}^n a_j e^{t_j z}$ obtained from (10) (and (9) too), is apparent from the determinant of $[\sum_{j=0}^n a_j e^{t_j z}]_{z=0}^{(i)}$, $i = 0(1)n$, which is the transposed Vandermondian $\|t_j^i\|$, $i, j = 0(1)n$.

CONCLUSION

In seeking new interpolation series, we have noted the heuristic value of employing different expansions of the same operational function (here it was $(1 + \Delta)^z$ or e^{Dz}), even though at each step (here for $z = i$ or for the i th derivative) those two different expansions of that same operational function become identical (here the same polynomials in Δ or D), which preserves the interpolation property. This suggests further applications of this principle to other expansions of $E^z = (1 + \Delta)^z = e^{Dz}$, in terms of different functions of the operators E , Δ or D , those in terms of polynomials (e.g., Legendre, Chebyshev, etc.) producing series that give either $f(i)$ for $z = i$, or $f^{(i)}(0)$ when differentiated i times and z set equal to zero, and those in terms of transcendental functions presenting interesting problems regarding their interpretation, convergence and utility, even when the series produced is not interpolatory.

REFERENCES

1. J. W. Layman, A finite difference exponential approximation method. *Math. Comp.* **18**, 113-118 (1964).
2. J. W. Layman, Mixed algebraic-exponential interpolation using finite differences. *Math. Comp.* **21**, 498-500 (1967).
3. J. W. Layman, Expansion of analytic functions in exponential polynomials. *Proc. Am. Math. Soc.* **22**, 519-522 (1969).
4. J. W. Layman, Convergence rates for exponential interpolation series. *J. Approx. Theory* **9**, 367-372 (1973).
5. A. V. Lotockii, A certain interpolation series (Russian), *Ivanov Gos. Ped. Inst. Učen. Zap.* **31**, 19-28 (1963); (Information from *Math. Rev.* **46**, p. 1621, no. 9359 (1973)).